




ESP Kick-Off Workshop Project Plan Presentation

Global Simulation of Plasma Microturbulence at the Petascale & Beyond

PI: William Tang (PPPL)

Presenter: Tim Williams (ANL)

October 18-19, 2010



Project Overview

Kinetic simulation of turbulent energy transport in magnetic fusion devices

- GTC-P
 - Large-aspect-ratio equilibrium magnetic geometry
 - Circular cross-section
- GTS
 - Extreme D-shape cross sections of existing tokamak fusion devices
 - Initial magnetic equilibrium set from experimental parameters
- Mira enables physical ITER scale simulations previously not possible on Intrepid
- Validated simulations & associated understanding of gyro-Bohm confinement would be major milestone for ITER
 - Insensitivity to increased system size (ITER-scale) associated with Gyro-Bohm confinement is a major advantage over unfavorable Bohm-like scaling observed in most present-generation tokamaks

Scientific Field: Fusion

Codes: GTS, GTC-P

Computational Approach, Numerical Methods

Gyrokinetic plasma description efficient for microturbulence regime

- Solve gyrophase-averaged Vlasov-Poisson
- Removes the fast cyclotronic time scale while keeping all non-local effects of gyration (particles are charged rings)

Particle-in-cell plasma simulation

- Charge of discrete simulation particles deposited to grid
- Self-consistent (electrostatic) field from solution of Poisson equation
 - PETSc solver (GMRES + Newton pre-conditioner)
- Particles move under force interpolated from field
- GTC-P: adiabatic electrons
- GTS: full electron dynamics

Parallelism and Existing Implementation

MPI parallelism:


- 1D domain decomposition in toroidal direction (long way around)
- GTS: Particles distributed within subdomain
- GTC-P: 2nd dimension domain decomposition (radial)


OpenMP parallelism

- Loop-level directives
- I/O: ADIOS
 - Only for checkpoint-restart at the moment

Parallelism and Existing Implementation (2)

Current Performance/Scalability





Library and Tool Dependencies

Libraries

- PETSc (solver for Poisson equation)
- ADIOS (parallel I/O)
- SPRNG 2.0 (random number generators)
- PSPLINE (spline and hermite interpolation tools)

Tools

VisIt

Anticipated Modifications for Blue Gene/Q

- Node-level parallelism
 - Increase ratio of OpenMP parallelism to MPI processes
- Processor-specific optimization at core level
- Port GTC-P optimizations to GTS
 - 2D domain decomposition
- Performance and scaling needed to run proposed problem on Mira:
 - ITER minor radius 4x current experiments such as DIII-D
 - 600 million grid cells
 - 50 billion particles
 - Need O(4x) speed of Intrepid for reasonable wallclock turnaround
 - Scale out to Intrepid cores = 640,000 Mira cores
 - About 320,000 Mira cores

Plan for Next 6 Months Effort

- Help find and hire a project postdoc
- On Intrepid:
 - Measure GTC-P performance w.r.t. increased OpenMP and decreased MPI
 - Project to estimate Mira performance
 - Simultaneously start work on optimizing multithread charge accumulation
 - Diagnose GTC-P load imbalance beyond 32K cores
 - Need to balance grid work (radial decomposition) with particle work
 - Manifested in local force calculation
 - Port GTC-P optimizations to GTS

